

Viscosity Studies on Lithium Bromide in Water + Dimethyl Sulfoxide Mixtures at 278.15 K and 293.15 K

Maria M. Palaiologou,* Ioanna E. Molinou, and Nikos G. Tsierkezos

Physical Chemistry Laboratory, Department of Chemistry, University of Athens, P.O. Box 64004, 15710 Zografou, Athens, Greece

Viscosities of dilute solutions of lithium bromide in binary mixtures of water + dimethyl sulfoxide (DMSO) have been measured in the composition range 0–85 mass % DMSO at 278.15 K and 0–100 mass % DMSO at 293.15 K. The viscosity data have been analyzed by the Jones–Dole equation. The viscosity B coefficients of LiBr are positive at both temperatures in the whole composition range of the mixed solvent. The ion–solvent interactions have been discussed.

1. Introduction

Viscosimetric measurements of electrolyte solutions have been widely used in order to obtain information regarding solute–solvent interactions. While ion–solvent interactions have been studied in water and some organic solvents, less consideration has been given to such effects in solvent mixtures. For such studies, the solvent mixture of water + DMSO has special importance because of the wide use of DMSO and its aqueous solutions as solvents and reaction media.¹ Also, this solvent mixture is of extreme interest from the point of view of their structural properties. Extensive studies of the physical properties of the binary system water (1) + DMSO (2) have been made to investigate the changes which occur in the molecular structure and motion of the liquid as a function of the composition. At ambient pressure, DMSO exists in the liquid state over a wide temperature range ($T_f = 18\text{ }^\circ\text{C}$ and $T_b = 189\text{ }^\circ\text{C}$) and is completely miscible with water. The high boiling temperature has been attributed to the large dipole moment of the DMSO molecule ($\mu = 3.90\text{ D}$ at $25\text{ }^\circ\text{C}$). In the strongly polar S=O bond, oxygen is the most electronegative site of DMSO. Dipole or ionic interactions are responsible for self-association DMSO molecules. Acting as a proton acceptor, it can alter the structure of water. From thermodynamic considerations, Lindberg² has argued that the water–DMSO hydrogen bond is stronger than the water–water hydrogen bond. Results of neutron inelastic scattering (NIS) and X-ray³ diffraction techniques show that small quantities of DMSO ($x_2 \leq 0.1$) added to water produce a cooperative ordering of water molecules, increasing the long-range structure of water. At higher concentrations of DMSO, the NIS, infrared spectroscopy, and X-ray results show that the water structure is broken down by the increased formation of hydrogen-bonded DMSO–water complexes.

On the other hand, infrared spectroscopy⁴ as well as density⁵ measurements, shows that small quantities of DMSO have little effect on the water structure. Other methods^{6,7} even lead to the conclusion that small amounts of DMSO act as a “structure breaker” in water. The dielectric spectra⁸ of water (1) + DMSO (2) mixtures showed two maxima (one each on the water rich region and DMSO rich region) in the distribution of relaxation times against concentration and a minimum around $x_2 = 0.33$, indicating more homogeneity around this mole fraction.

Table 1. Densities (ρ) and Viscosities (η) of Water (1) + DMSO (2) Mixtures at 278.15 K and 293.15 K

100w ₂	T = 278.15 K		T = 293.15 K	
	$\rho/\text{g}\cdot\text{cm}^{-3}$	$\eta/\text{mPa}\cdot\text{s}$	$\rho/\text{g}\cdot\text{cm}^{-3}$	$\eta/\text{mPa}\cdot\text{s}$
0.00	0.999 96	1.524	0.998 20	0.999
2.50	1.003 32	1.589	1.001 32	1.053
5.00	1.006 89	1.727	1.004 52	1.112
10.00	1.014 31	1.948	1.011 20	1.264
14.97	1.022 00	2.178	1.018 20	1.402
18.15	1.027 27	2.354	1.022 83	1.515
20.00	1.030 38	2.463	1.025 53	1.559
30.00	1.047 50	3.203	1.040 96	2.009
40.00	1.065 05	4.230	1.056 62	2.589
42.75	1.070 15	4.625	1.061 69	2.743
50.00	1.082 53	5.570	1.072 40	3.319
65.00	1.104 44	7.330	1.092 33	4.296
70.00	1.109 25	7.326	1.096 70	4.385
78.00	1.115 47	6.666	1.102 16	4.091
85.00	1.117 61	5.450	1.103 81	3.498
89.00	1.117 88	4.904	1.103 71	3.211
93.00	1.117 48	4.035	1.102 91	2.743
96.00	1.117 02	3.630	1.102 12	2.514
100.00			1.100 41	2.194
			1.100 40 ^a	2.216 ^a

^a Reference 31.




Figure 1. Viscosity of water (1) + DMSO (2) mixtures: ■, 278.15 K; ●, 293.15 K.

The existence of stoichiometrically well-defined hydrogen-bonded DMSO–water aggregates is suggested thereby.

Table 2. Experimental Densities (ρ) and Viscosities (η) of LiBr in Water (1) + DMSO (2) Mixtures at 278.15 K and 293.15 K

$c/\text{mol}\cdot\text{dm}^{-3}$		$\rho/\text{g}\cdot\text{cm}^{-3}$		$\eta/\text{mPa}\cdot\text{s}$		$c/\text{mol}\cdot\text{dm}^{-3}$		$\rho/\text{g}\cdot\text{cm}^{-3}$		$\eta/\text{mPa}\cdot\text{s}$	
278.15 K						293.15 K					
100 w ₂ = 0.00						100 w ₂ = 0.00					
0.0098	1.000 60	1.500	0.0453	1.002 86	1.506	0.0146	0.999 14	1.001	0.0453	1.001 07	1.005
0.0146	1.000 91	1.501	0.0543	1.003 45	1.507	0.0198	0.999 46	1.002	0.0542	1.001 63	1.006
0.0199	1.001 25	1.502	0.0802	1.005 07	1.511	0.0279	0.999 97	1.003	0.0801	1.003 25	1.009
0.0280	1.001 75	1.503	0.0996	1.006 36	1.514	0.349	1.000 43	1.004	0.1000	1.004 51	1.012
0.0350	1.002 20	1.504	0.1222	1.007 81	1.516						
100 w ₂ = 2.50						100 w ₂ = 2.50					
0.0149	1.004 28	1.592	0.0558	1.006 90	1.599	0.0149	1.002 28	1.055	0.0557	1.004 82	1.060
0.0235	1.004 83	1.594	0.0808	1.008 48	1.603	0.0235	1.002 81	1.056	0.0812	1.006 38	1.063
0.0284	1.005 24	1.595	0.1018	1.009 82	1.606	0.0283	1.003 22	1.057	0.1016	1.007 75	1.066
0.0350	1.005 65	1.596	0.1176	1.010 84	1.609	0.0349	1.003 57	1.058	0.1173	1.008 72	1.068
0.0456	1.006 24	1.598				0.0455	1.004 20	1.059			
100 w ₂ = 5.00						100 w ₂ = 5.00					
0.0186	1.008 07	1.718	0.0749	1.011 63	1.730	0.0135	1.005 38	1.115	0.0415	1.007 10	1.119
0.0257	1.008 53	1.720	0.0955	1.012 95	1.733	0.0178	1.005 63	1.116	0.0470	1.007 46	1.120
0.0316	1.008 92	1.721	0.1207	1.014 49	1.738	0.0200	1.005 77	1.116	0.0584	1.008 09	1.122
0.0484	1.009 97	1.724				0.0308	1.006 45	1.117	0.1246	1.012 28	1.130
						0.0386	1.006 92	1.118			
100 w ₂ = 10.00						100 w ₂ = 10.00					
0.0050	1.014 64	1.949	0.0320	1.016 32	1.954	0.0104	1.011 85	1.267	0.0363	1.013 46	1.273
0.0104	1.014 96	1.950	0.0364	1.016 59	1.955	0.0137	1.012 06	1.268	0.0443	1.013 94	1.275
0.0137	1.015 19	1.951	0.0445	1.017 12	1.957	0.0171	1.012 27	1.269	0.0506	1.014 33	1.276
0.0172	1.015 41	1.952	0.0508	1.017 49	1.958	0.0264	1.012 82	1.271	0.1059	1.017 71	1.285
0.0265	1.015 99	1.953	0.1062	1.020 90	1.969	0.0320	1.013 19	1.272			
100 w ₂ = 15.00						100 w ₂ = 15.00					
0.0154	1.022 96	2.181	0.0809	1.026 41	2.198	0.0153	1.019 02	1.405	0.0806	1.022 48	1.418
0.0207	1.023 17	2.184	0.1025	1.027 57	2.204	0.0206	1.019 60	1.407	0.1021	1.023 67	1.421
0.0263	1.023 49	2.185	0.1526	1.030 24	2.214	0.0262	1.019 60	1.408	0.1520	1.026 41	1.429
0.0518	1.024 87	1.192				0.0516	1.020 99	1.412			
100 w ₂ = 18.15						100 w ₂ = 18.15					
0.0164	1.028 11	2.361	0.0846	1.032 10	2.381	0.0207	1.024 11	1.522	0.1020	1.028 77	1.541
0.0214	1.028 52	2.363	0.1024	1.033 13	2.388	0.0273	1.024 58	1.524	0.1214	1.029 77	1.544
0.0274	1.028 86	2.365	0.1219	1.034 17	2.393	0.0521	1.025 87	1.530	0.2003	1.034 36	1.562
0.0523	1.030 31	2.373									
100 w ₂ = 20.00						100 w ₂ = 20.00					
0.0181	1.031 48	2.470	0.0425	1.033 03	2.480	0.0063	1.025 95	1.562	0.0423	1.028 17	1.572
0.0199	1.031 56	2.471	0.0481	1.033 38	2.482	0.0180	1.026 66	1.565	0.0479	1.028 55	1.573
0.0301	1.032 17	2.474	0.1015	1.036 63	2.502	0.0198	1.026 80	1.566	0.0605	1.029 31	1.577
0.0396	1.032 78	2.478	0.1532	1.039 93	2.521	0.0300	1.027 40	1.568	0.1011	1.031 82	1.588
						0.0394	1.028 02	1.571			
100 w ₂ = 30.00						100 w ₂ = 30.00					
0.0150	1.048 33	3.215	0.1017	1.053 12	3.263	0.0150	1.041 89	2.015	0.1011	1.046 62	2.049
0.0207	1.048 58	3.217	0.1212	1.054 19	3.275	0.0206	1.042 16	2.018	0.1204	1.047 73	2.058
0.0504	1.050 32	3.236	0.1524	1.055 96	3.296	0.0501	1.043 84	2.030	0.1515	1.049 53	2.070
0.0818	1.052 02	3.255				0.0813	1.045 56	2.042			
100 w ₂ = 40.00						100 w ₂ = 40.00					
0.0120	1.065 77	4.249	0.0463	1.067 84	4.284	0.0202	1.057 91	2.607	0.0570	1.060 18	2.632
0.0154	1.065 98	4.252	0.0575	1.068 73	4.302	0.0289	1.058 45	2.612	0.0838	1.061 80	2.651
0.0203	1.066 28	4.256	0.0845	1.070 17	4.327	0.0364	1.058 90	2.617	0.1037	1.063 00	2.664
0.0292	1.066 84	4.266	0.1045	1.071 35	4.345	0.0459	1.059 49	2.625	0.1299	1.064 68	2.681
0.0367	1.067 43	4.276	0.1309	1.072 94	4.376						
100 w ₂ = 42.75						100 w ₂ = 42.75					
0.0062	1.070 40	4.643	0.1139	1.077 25	4.805	0.0061	1.062 00	2.753	0.0970	1.066 52	2.830
0.0164	1.071 33	4.660	0.1481	1.079 15	4.839	0.0162	1.062 44	2.761	0.1139	1.068 67	2.850
0.0490	1.073 07	4.709	0.1930	1.082 45	4.884	0.0221	1.063 18	2.767	0.1469	1.070 63	2.872
0.0897	1.073 80	4.755				0.0488	1.064 70	2.789	0.1892	1.073 96	2.914
100 w ₂ = 49.95						100 w ₂ = 49.95					
0.0142	1.083 43	5.608	0.0594	1.086 24	5.702	0.0106	1.073 00	3.332	0.0487	1.075 39	3.375
0.0194	1.083 80	5.619	0.0855	1.087 77	5.748	0.0141	1.073 28	3.337	0.0589	1.076 06	3.388
0.0259	1.084 12	5.632	0.1071	1.089 09	5.791	0.0193	1.073 58	3.342	0.0847	1.077 60	3.414
0.0315	1.084 45	5.644	0.1286	1.090 32	5.828	0.0258	1.073 98	3.350	0.1061	1.078 92	3.438
0.0371	1.084 75	5.654	0.1616	1.092 33	5.896	0.0313	1.074 32	3.356	0.1283	1.080 22	3.460
0.0492	1.085 51	5.674				0.0367	1.074 64	3.362			
100 w ₂ = 65.00						100 w ₂ = 65.00					
0.0152	1.105 19	7.383	0.1042	1.109 84	7.709	0.0151	1.093 13	4.327	0.1030	1.097 83	4.488
0.0200	1.105 47	7.399	0.1254	1.111 00	7.769	0.0198	1.093 40	4.336	0.1241	1.099 01	4.526
0.0249	1.105 66	7.416	0.1572	1.112 61	7.875	0.0247	1.093 69	4.347	0.1555	1.100 66	4.584
0.0509	1.107 07	7.511	0.2095	1.115 39	8.077	0.0504	1.095 06	4.390	0.2073	1.103 43	4.677
0.0814	1.108 74	7.628	0.2574	1.117 59	8.240	0.0805	1.096 70	4.447	0.2546	1.105 65	4.758
100 w ₂ = 84.98						100 w ₂ = 84.98					
0.0156	1.118 52	5.516	0.0827	1.122 54	5.833	0.0154	1.104 78	3.539	0.1029	1.110 18	3.773
0.0209	1.118 89	5.545	0.1042	1.123 91	5.937	0.0206	1.105 40	3.558	0.1245	1.111 49	3.842
0.0259	1.119 16	5.577	0.1581	1.127 17	6.212	0.0256	1.105 42	3.564	0.1562	1.113 42	3.934
0.0520	1.120 79	5.687	0.2074	1.130 11	6.467	0.0515	1.107 00	3.642	0.2048	1.116 40	4.075
						0.0817	1.108 84	3.723			
100 w ₂ = 100.00						100 w ₂ = 100.00					
						0.0112	1.101 23	2.216	0.0487	1.103 97	2.297
						0.0159	1.101 58	2.226	0.0596	1.104 76	2.320
						0.0221	1.102 02	2.239	0.0690	1.105 45	2.342
						0.0303	1.102 62	2.257	0.1083	1.108 31	2.430
						0.0379	1.103 18	2.273			

Table 3. Viscosity *A* and *B* Coefficients of the Jones–Dole Equation (Eq 1) for LiBr in Water (1) + DMSO (2) Mixtures at 278.15 and 293.15 K

100 <i>w</i> ₂	<i>A</i> /(dm ³ ·mol ⁻¹) ^{1/2}		<i>B</i> /dm ³ ·mol ⁻¹	
	<i>T</i> = 278.15 K	<i>T</i> = 293.15 K	<i>T</i> = 278.15 K	<i>T</i> = 293.15 K
0.00	0.0012 ± 0.0001	0.0053 ± 0.0005	0.093 ± 0.002	0.104 ± 0.002
2.50	0.0040 ± 0.0004	0.0075 ± 0.0008	0.092 ± 0.002	0.102 ± 0.002
5.00	0.0112 ± 0.0011	0.0107 ± 0.0011	0.089 ± 0.002	0.096 ± 0.002
10.00	0.0006 ± 0.0001	0.0209 ± 0.0021	0.098 ± 0.002	0.098 ± 0.002
15.00	0.0030 ± 0.0003	0.0094 ± 0.0009	0.102 ± 0.002	0.105 ± 0.002
18.15	0.0110 ± 0.0011	0.0166 ± 0.0017	0.104 ± 0.002	0.114 ± 0.002
20.00	-0.0003 ± 0.0001	0.0050 ± 0.0005	0.158 ± 0.003	0.163 ± 0.003
30.00	0.0070 ± 0.0007	0.0020 ± 0.0002	0.170 ± 0.003	0.196 ± 0.004
40.00	0.0150 ± 0.0015	0.0142 ± 0.0014	0.219 ± 0.004	0.234 ± 0.004
42.75	0.0310 ± 0.0031	0.0193 ± 0.0019	0.228 ± 0.004	0.276 ± 0.005
49.95	0.0189 ± 0.0019	0.0076 ± 0.0008	0.311 ± 0.006	0.312 ± 0.006
65.00	0.0007 ± 0.0001	0.0091 ± 0.0009	0.483 ± 0.010	0.406 ± 0.008
85.00	-0.0131 ± 0.0013	-0.0053 ± 0.0005	0.917 ± 0.018	0.807 ± 0.016
100.00		-0.0151 ± 0.0015		1.036 ± 0.021
				0.907 ± 0.003 ^a

^a Value at 298.15 K.⁴⁰

The studies indicate a strong interaction between DMSO and water, and the general conclusion is that in the mole fraction range $x_2 = 0.3$ to 0.4 the interactions due to hydrogen bonds are at a maximum. When plotted against concentration, properties of the mixtures such as density,⁹ viscosity,⁹ spin lattice relaxation rates,¹⁰ enthalpy of mixing,¹¹ and surface tension¹² pass through minima or maxima at about $x_2 = 0.35$. There is some disagreement as to whether actual hydrogen-bonded complexes are formed,^{3,13,14} what the water to DMSO ratio is in the complex,¹⁵ and how long such complexes exist.^{10,16} For example, measurements of the neutron spin–lattice relaxation time in the water + DMSO system have suggested that the reorientation of the DMSO·3H₂O complex is the dominant motion causing relaxation.¹⁵ Also, from phase diagram studies, it is concluded that a complex of one DMSO and three water molecules is formed.¹⁷

Other dielectric studies have also shown that a DMSO molecule reorients together with two water molecules.¹⁶ This is in agreement with enthalpy of mixing,¹¹ density,⁹ and viscosity⁹ data, where the results also indicate a DMSO·2H₂O complex.

From a phase diagram of the supercooled aqueous solutions¹⁸ of DMSO it is suggested that a stable complex of DMSO·2H₂O exists the near glass transition temperature (T_g) in DMSO.

The existence of strongly hydrogen bonded DMSO·2H₂O aggregates has been proved through molecular dynamical simulations of these solvent mixtures.¹⁹ The behavior of the principal relaxation time of the mixture shows a maximum at a value corresponding to a mixture with 30 mass % water.²⁰

A survey of the literature shows that ionic viscosity *B* coefficients have been investigated for electrolyte solutions in water,^{21–23} in DMSO,^{24–26} and in water + DMSO mixtures.^{27–30} This work involves the study of the *B* coefficient of lithium bromide in water + DMSO mixtures in the whole mole fraction region at 278.15 K and 293.15 K.

2. Experimental Section

Dimethyl sulfoxide (Riedel–de Hën, 99.9%) with density 1.100 41 g·cm⁻³ and refractive index 1.4795 at 293.15 K was used without further purification. These physical constants are in agreement with the literature values $\rho = 1.100 401$ g·cm⁻³ and $n_D = 1.4793$.³¹ Water was deionized by an ion-exchange resin and doubly distilled. The specific

conductance was found to be better than $0.50 \mu\text{S}\cdot\text{cm}^{-1}$ at 293.15 K. Anhydrous lithium bromide (Fluka 99.0%) was dried before use at 413.15 K for 24 h. All solutions were prepared by mass from a stock solution. The weighings were uncertain to less than ± 0.0001 g (salt) and ± 0.01 g (solvent mixtures). Molar concentrations were obtained from molality and density values with an uncertainty ± 0.0001 mol·dm⁻³.

Densities (ρ) were measured with an Anton Paar (DMA 58) microcomputer-controlled precision densimeter, with a build-in solid state thermostat. The precision of the sample thermostat was ± 0.01 K. The DMA cell was calibrated with air and doubly distilled water at atmospheric pressure. The uncertainty in the density was $\pm 0.000 01$ g·cm⁻³.

The viscosities have been measured in the concentration range 0.0010–0.1500 mol·dm⁻³ with a viscosity measuring unit (Schott Geräte AVS 310) equipped with an Ubbelohde capillary viscometer. The viscometer was calibrated with doubly distilled water. The time measurement tolerance was $\pm 0.005\%$, and the display accuracy was ± 0.01 s. The temperature was maintained constant within ± 0.03 K. The viscosities (η) were averaged from four readings. The uncertainty in the viscosity measurements was ± 0.001 mPa·s.

3. Results and Discussion

Densities and viscosities of pure water, DMSO, and water + DMSO mixtures are reported in Table 1. Inspection of density values shows that the density passes through a maximum between 80 and 90 mass % DMSO for both temperatures. The viscosity–concentration curves, plotted in Figure 1, exhibit very pronounced maxima, which occur at a concentration corresponding to the formula DMSO/2H₂O (100 *w*₂ = 68). These maxima in the viscosity water + DMSO system can be attributed to the strong interaction of DMSO with water. These results agree well with the literature values.^{9, 32–34}

Densities and viscosities of LiBr in water + DMSO mixtures are reported in Table 2.

The viscosities of electrolytic solutions were analyzed using the Jones–Dole³⁵ equation:

$$\eta_r = 1 + Ac^{1/2} + Bc \quad (1)$$

where η_r is the relative viscosity of the solution, the *A* coefficient is a measure of ion–ion interactions, and the *B*




Figure 2. Viscosity B coefficients for LiBr in water (1) + DMSO (2) mixtures: ■, 278.15 K; ●, 293.15 K.

coefficient depends on the specific electrolyte and temperature and consequently is a measure of ion–solvent interactions.

The viscosity A and B coefficients have been calculated as the intercept and the slope, respectively, from plots of $(\eta/\eta_0 - 1)/c^{1/2}$ against $c^{1/2}$. The A and B values derived from eq 1 are given with their standard deviations in Table 3.

It is seen that the viscosity A coefficient is negative in some cases, but this has no physical significance.³⁶

The values of the B coefficient for both temperatures are presented graphically in Figure 2. The B coefficient is positive throughout all solvent mixtures, indicating a structure making behavior of LiBr. In pure water the Li⁺ ion is considered to be a structure making ion whereas the Br⁻ ion is considered to be a structure breaking ion.²³

As is shown in Figure 2, the change of the B coefficient at both temperatures is very small up to $100w_2 \sim 18.15$, within the experimental uncertainty, with a small decrease from water to $100w_2 = 10$. This change of the B values in the water rich region may be due to the weakening of the ion–solvent interactions, because of the domination of the solvent–solvent interactions between unlike solvent molecules. Above $100w_2 = 18.15$ there is a marked increase in the B values. This increase may be attributed to the stronger solvation of Li⁺ by DMSO molecules because of their greater basicity (donor number = 29.8 kJ·mol⁻¹).³⁷ On the other hand, the Br⁻ ion exhibits weaker solvation because the positive end of the DMSO dipole is sterically hindered. The preference solvation of Li⁺ and the small solvation of Br⁻ by DMSO are also confirmed from the free energies of transfer³⁸ of ions from water to water + DMSO mixtures and from the limiting ionic conductivities³⁹ in DMSO. Therefore, the structure making behavior of the salt can be attributed mainly to Li⁺.

In conclusion, the B coefficient is positive throughout the solvent mixtures for both temperatures, showing a structure maker behavior of LiBr. The thermal coefficient (dB/dT) is positive up to $100w_2 = 50$, and above this composition turns to negative values. This change may be due to the strongest solvent–solvent interactions in the lower temperature. Though at higher concentration of DMSO the solvent–solvent interactions are less strong, the effect of long-range ordering of the solvent around the ions decreases with the increasing thermal energy.²³

Literature Cited

- Fuchs, R.; McCrary, G. E.; Bloomfield, J. J. Mechanisms of Nucleophilic Displacement in Aqueous Dimethyl Sulfoxide Solutions. *J. Am. Chem. Soc.* **1961**, *83*, 4281–4284.
- Lindberg, J. J. Dimethyl sulfoxide, structure and physicochemical properties. *Suom. Kemistiseuran Tied.* **1961**, *70*, 33–39.
- Safford, G. J.; Schaffer, P. C.; Leung, P. S.; Doebbler, G. F.; Brady, G. W.; Lyden, E. F. X. Neutron Inelastic Scattering and X-ray Studies of Aqueous Solutions of Dimethylsulphoxide and Dimethylsulphone. *J. Chem. Phys.* **1969**, *50*, 2140–2159.
- Brink, G.; Falk, M. The effect of dimethyl sulphoxide on the structure of water. *J. Mol. Struct.* **1970**, *5*, 27–30.
- McDonald, D. D.; Smith, M. D.; Hyne, J. B. The Influence of Sulfoxides and Sulfones on the Temperature of Maximum Density of Water. *Can. J. Chem.* **1971**, *49*, 2817–2821.
- De Visser, C.; Heuvelsland, M. J. M.; Dunn, L. A.; Somsen, G. Some Properties of Binary Aqueous Liquid Mixtures. Apparent Molar Volumes and Heat Capacities at 298.15 K Over the Whole Mole Fraction Range. *J. Chem. Soc., Faraday Trans 1* **1978**, *74*, 1159–1169.
- Petrella, G.; Petrella, M.; Gastagnolo, M.; Dell'Atti, A.; De Giglio, A. Solute–Solvent Interactions in Water-Rich Mixtures. Ionic Conductances in Water-Dimethyl sulfoxide Mixtures at 25 °C. *J. Solution Chem.* **1981**, *10*, 129–138.
- Kaatz, U.; Pottel, R.; Schafer, M. Dielectric Spectrum of Dimethyl sulfoxide/Water Mixtures as a Function of Composition. *J. Phys. Chem.* **1989**, *93*, 5623–5627.
- Cowie, J. M. G.; Toporowski, P. M. Association in the Binary Liquid System Dimethyl Sulphoxide – Water. *Can. J. Chem.* **1961**, *39*, 2240–2243.
- Packer, K. J.; Tomlinson, J. D. Nuclear Spin Relaxation and Self-Diffusion in the Binary System, Dimethylsulphoxide (DMSO)+ Water. *J. Chem. Soc., Faraday Trans.* **1971**, *67*, 1302–1314.
- Fox, M. F.; Whittingham, K. P. Component Interactions in Aqueous Dimethyl Sulphoxide. *J. Chem. Soc., Faraday Trans.* **1975**, *71*, 1407–1412.
- Luzar, A. The Contribution of Hydrogen Bonds to Bulk and Surface Thermodynamic Properties of Dimethyl sulfoxide-Water Mixtures. *J. Chem. Phys.* **1989**, *91*, 3603–3613.
- Lindberg, J. J.; Majani, C. Notes on Solvate Formation in Dimethyl Sulphoxide-Water Mixtures. *Acta Chem. Scand.* **1963**, *17* (5), 1477–1478.
- Madigosky, W. M.; Warfield, R. W. Ultrasonic measurements and liquid structure of DMSO-water mixture. *J. Chem. Phys.* **1983**, *78*, 1912–1916.
- Glaser, J. A. Deuteron Magnetic Relaxation Studies on the Solution Properties of Some Denaturing Agents and Surfactants. *J. Am. Chem. Soc.* **1970**, *92*, 372–375.
- Higashigaki, Y.; Christensen, D. H.; Wang, C. H. Studies of the Reorientational Motion and Intermolecular Interaction of Dimethyl Sulfoxide in Water by Depolarized Rayleigh Scattering. *J. Phys. Chem.* **1981**, *85*, 2531–2535.
- Rasmussen, D. H.; Mackenzie, A. P. Phase Diagram for the System Water-Dimethylsulphoxide. *Nature* **1968**, *220*, 1315–1317.
- Murthy, S. S. N. Phase Behavior of the Supercooled Aqueous Solutions of Dimethyl Sulfoxide, Ethylene Glycol, and Methanol as Seen by Dielectric Spectroscopy. *J. Phys. Chem. B* **1997**, *101*, 6043–6049.
- Vaisman, I. I.; Berkowitz, M. L. Local Structural Order and Molecular Associations in Water-DMSO Mixtures. Molecular Dynamics Study. *J. Am. Chem. Soc.* **1992**, *114*, 7889–7896.
- Puranik, S. M.; Kumbharhane, A. C.; Mehrotra, S. C. Dielectric Study of Dimethyl Sulfoxide-Water Mixtures Using the Time-domain Technique. *J. Chem. Soc., Faraday Trans.* **1992**, *88*, 433–435.
- Desnoyers, J. E.; Perron, G. The Viscosity of Aqueous Solutions of Alkali and Tetraalkylammonium Halides at 25 °C. *J. Solution Chem.* **1972**, *1*, 199–212.
- Sacco, A.; Giglio, A.; Dell'Atti, A.; Petrella, M. Ionic B Coefficients in Water at 30, 40, and 50 °C. *J. Chem. Soc., Faraday Trans. 1* **1981**, *77*, 2693–2693.
- Kaminsky, M. Ion–Solvent Interaction and the Viscosity of Strong-Electrolyte Solutions. *Discuss. Faraday Soc.* **1957**, *24*, 171–179.
- Bicknell, R. T. M.; Lawrence, K. G.; Seeley, M.-A.; Feakins, D.; Werblan, L. Relative Viscosities of Solutions of Sodium and Potassium Bromides and Iodides in Dimethyl Sulphoxide at 25, 35 and 45 °C. *J. Chem. Soc., Faraday Trans.* **1976**, *72*, 307–313.
- Bicknell, R. T. M.; Lawrence, K. G.; Feakins, D. Ionic Viscosity B Coefficients in Dimethyl Sulphoxide at 25, 35, and 45 °C. *J. Chem. Soc., Faraday Trans. 1* **1980**, *76*, 637–647.
- Lawrence, K. G.; Sacco, A. Separation of Viscosity B Coefficients into Ionic Contributions. *J. Chem. Soc., Faraday Trans. 1* **1983**, *79*, 615–624.
- Sacco, A.; De Giglio, A.; Castagnolo, M.; Dell'Atti, A. Ion–Solvent Interactions in water-Rich Binary Mixtures. Part 3. Viscometric Behaviour of NaCl, NaBr, NaI and KCl in Water + Dimethylsulphoxide Mixtures at 25 °C and 35 °C. *Z. Phys. Chem. Neue Folge* **1986**, *150*, 105–112.
- Osborne, D. W.; O'Brien, J. F. Viscosities of Solutions of Lithium Chloride in Water-Dimethyl Sulfoxide Mixtures at 25, 35, and 45 °C. *J. Chem. Eng. Data* **1986**, *31*, 317–320.

- (29) Jauhar, S. P.; Guraya, P. S.; Parmar, J. S.; Narula, S. P. Physico-Chemical Studies in Mixed Solvents. Part II. Viscosity Measurements of Some 1: 1 Electrolyte Solutions in Dimethylsulphoxide-Water Mixtures at 25 °C. *Z. Phys. Chem. Neue Folge* **1985**, *146*, 9–13.
- (30) Campbell, A. N. Electrolytes in mixed solvents: indium trichloride in water-dimethyl sulfoxide mixtures. *Can. J. Chem.* **1978**, *56*, 355–358.
- (31) Bicknell, R. T. M.; Davies, D. B.; Lawrence, K. G. Density, Refractive Index, Viscosity and ¹H Nuclear Magnetic Resonance Measurements of Dimethyl Sulphoxide at 2 °C Intervals in the Range 20–60 °C. *J. Chem. Soc., Faraday Trans. 1* **1982**, *78*, 1595–1601.
- (32) Schichman, S. A.; Amey, R. L. Viscosity and Local Liquid Structure in Dimethyl Sulfoxide-Water Mixtures. *J. Phys. Chem.* **1971**, *75*, 98–102.
- (33) Fort, R. J.; Moore, W. R. Viscosities of Binary Liquid Mixtures. *J. Chem. Soc., Faraday Trans. 1* **1966**, *62*, 1112–1119.
- (34) Campbell, A. N.; Kartzwark, E. M. Conductivities of Lanthanum Chloride in Water – Dimethyl Sulfoxide Mixtures at 25 °C. *J. Chem. Eng. Data* **1984**, *29*, 168–171.
- (35) Jones, G.; Dole, M. The Viscosity of Aqueous Solutions of Strong Electrolytes With Special Reference to Barium Chloride. *J. Am. Chem. Soc.* **1929**, *51*, 2950–2964.
- (36) Harned, H. S.; Owen, B. *Physical Chemistry of Electrolytic Solutions*; Reinhold Publishing Corporation: New York, 1943.
- (37) Gutmann, V. *The Donor–Acceptor Approach to Molecular Interactions*; Plenum Press: New York, 1978.
- (38) Cox, B. G.; Natarajan, R. Thermodynamic Properties for Transfer of Electrolytes from Water to Dimethylsulphoxide and to Dimethylsulphoxide + Water Mixtures. *J. Chem. Soc., Faraday Trans. 1979*, *75*, 1780–1787.
- (39) Kale, K. M.; Zana, R. Partial Molal Volumes of Ions in Organic Solvents from Ultrasonic Vibration Potential and Density Measurements. III. Dimethyl Sulfoxide. *J. Solution Chem.* **1977**, *6*, 733–746.

Received for review April 5, 2002. Accepted July 10, 2002. This work was financially supported by the Special Research Account of the National and Kapodistrian University of Athens.

JE020063S