

A Multireference Coupled-Cluster Potential Energy Surface of Diazomethane, CH₂N₂

Ioannis S. K. Kerkines,[†] Petr Čársky,[‡] and Aristides Mavridis^{*,†}

Laboratory of Physical Chemistry, Department of Chemistry, National and Kapodistrian University of Athens, PO Box 64 004, 157 10 Zografou, Athens, Greece, and J. Heyrovský Institute of Physical Chemistry, Academy of Sciences of the Czech Republic, Dolejškova 3, 182 23 Prague 8, Czech Republic


Received: August 12, 2005; In Final Form: September 10, 2005

The intrinsically multireference dissociation of the C–N bond in ground-state diazomethane (CH₂N₂) at different angles has been studied with the multireference Brillouin–Wigner coupled-cluster singles and doubles (MRBWCCSD) method. The morphology of the calculated potential energy surface (PES) in C_s symmetry is similar to a multireference perturbational (CASPT3) PES. The MRBWCCSD/cc-pVTZ H₂C–N₂ dissociation energy with respect to the asymptotic CH₂(\tilde{a}^1A_1) + N₂(X¹Σ_g⁺) products is D_e = 35.9 kcal/mol, or a zero-point corrected D₀ = 21.4 kcal/mol with respect to the ground-state CH₂(\tilde{X}^3B_1) + N₂(X¹Σ_g⁺) fragments.

1. Introduction

Diazomethane (CH₂N₂) is a well-known organic compound that was discovered more than a century ago (1894).¹ Despite its toxicity and explosiveness, it has for a long time been used in organic synthesis as a means of very efficient and facile acid, alcohol, and phenol methylation.¹ Recent studies have also indicated its probable formation in the N₂-rich ice surfaces of Triton (a moon of Neptune) and Pluto.^{2,3}

The various isomers of CH₂N₂, mainly diazomethane, diazirine, and cyanamide, have been largely studied in the past, both experimentally and theoretically.^{4,5} Indeed, by 1962, the experimental geometrical structures of diazomethane and diazirine were already known.^{6,7} From a chemical point of view, diazomethane is one of the many molecules for which a single “conventional” Lewis structure cannot be written. In literature, it is usually referred to as a closed-shell “1–3 dipolar resonance hybrid” of the form^{8,9}



but, a singlet biradical form has also been proposed by Walsh and Goddard:¹⁰



a conjecture strongly opposed by Gerratt and co-workers in the light of spin-coupled calculations.¹¹ Instead, it is proposed that the in situ central nitrogen atom is in a “hypervalent” state with a valence number of five:




A somehow similar point of view has been put forth by Papakondylis and Mavridis⁴ who, with the help of multireference perturbation calculations proposed a similar “five-bond” picture

of the central N, without the need though of employing any virtual “hypervalent” state of N:




This bonding picture is based on the actual spectroscopic states of the fragments that participate in the formation of the \tilde{X}^1A_1 state of diazomethane, CH₂ + N₂. The in situ CH₂ finds itself in its \tilde{c}^1A_1 excited state, whereas the N₂ moiety remains in its ground X¹Σ_g⁺ state according to the following valence-bond Lewis (vbL) diagrams:



The two fragments are connected with a dative σ-bond stemming from one of the N₂ lone pairs to the empty CH₂ p_σ orbital, and a π-“back-transfer” conjugative interaction from the doubly occupied p_π orbital of CH₂ to the π system of N₂. The –N₂ triple bond remains practically intact as it lengthens by only ~0.035 Å as compared with the free N₂ bond length.⁴

The characteristics of the chemically important dissociation of diazomethane to the CH₂ and N₂ fragments



have caught the attention of the scientific community for more than half a century. It was early realized that this process had an energy barrier in C_{2v} symmetry, attributed to a Woodward–Hoffmann nonfavored dissociation.¹² As early as 1976, Lievin and Verhaegen pointed out the participation of the highly excited \tilde{c}^1A_1 state of CH₂ to the existing barrier and its involvement to the diazomethane equilibrium structure.^{12,13} Even though their calculations along the CH₂ + N₂ singlet potential energy surface (PES) were rather minimal, these authors suggested that the

* Corresponding author. E-mail: mavridis@chem.uoa.gr.

[†] National and Kapodistrian University of Athens.


[‡] Academy of Sciences of the Czech Republic.

TABLE 1: Energies, Bond Lengths, Angles, and Energy Gaps of the Four Lowest-Lying States of CH₂ at the (CASPT3, MRBWCCSD, MRCI, MRCI+Q and MRACPF)/cc-pVTZ Levels of Theory

	$E (E_h)$	$R_{C-H} (\text{Å})$	$\angle H-C-H (\text{deg})$	$T_e (\text{kcal/mol})$
		\tilde{X}^3B_1		
CASPT3	-39.074522	1.078	133.7	0
MRBWCCSD	-39.074928 ^a			0
MRCI	-39.074542	1.0787	133.3	0
MRCI+Q	-39.078511 ^b			0
MRACPF	-39.077245	1.0788	133.4	0
MRCI ^c	-39.140113	1.0751	133.66	0
expt ^d		1.07530	133.9308	0
		\tilde{a}^1A_1		
CASPT3	-39.057323	1.110	101.5	10.79
MRBWCCSD	-39.060832 ^a			8.85
MRCI	-39.058916	1.1109	101.6	9.81
MRCI+Q	-39.063085 ^b			9.68
MRACPF	-39.061759	1.1107	101.7	9.72
MRCI ^c	-39.125487	1.1061	102.20	9.178
expt		1.107 ± 0.002^e	102.4 ± 0.4^e	9.354^d
		\tilde{b}^1B_1		
CASPT3	-39.019908	1.074	142.6	34.27
MRBWCCSD	-39.019484 ^a			34.79
MRCI	-39.019619	1.076	141.8	34.46
MRCI+Q	-39.024768 ^b			33.72
MRACPF	-39.023148	1.075	142.5	33.95
MRCI ^c	-39.087831	1.0723	142.44	32.807
expt ^f		1.086	139.3	
expt ^g		1.053	140 ± 15	
		\tilde{c}^1A_1		
CASPT3	-38.972117	1.068	173.9	64.26
MRBWCCSD	-38.974976 ^a			62.72
MRCI	-38.976951	1.066	173.9	61.23
MRCI+Q	-38.982550 ^b			60.22
MRACPF	-38.980791	1.068	171.3	60.53
MRCI ^c	-39.046391	1.0657	171.92	58.811

^a At the CASPT3-optimized geometry. ^b At the MRCI-optimized geometry. ^c All-electron calculation/d-aug-cc-pCV6Z basis, ref 24, ^e Reference 25. ^f T_0 value converted to T_e using the zero-point energy difference from ref 26. ^d Reference 27. ^g Reference 28. ^h Reference 29.

formation of diazomethane can be thought of as a two-step barrierless process in C_s symmetry involving a perpendicular π -attack of N₂ to CH₂ (\tilde{a}^1A_1) followed by relaxation of the system to a C_{2v} geometry with the active participation of the CH₂ \tilde{c}^1A_1 state, and an opening of the $\angle HCH$ angle from ~ 102 to $\sim 126^\circ$.¹² This was verified and vividly exhibited in the complete active space third-order perturbation theory (CASPT3) PES by Papakondylis and Mavridis⁴ with a correlation-consistent basis set of triple- ζ quality, cc-pVTZ.¹⁴ The reason for the preferred barrierless approach of N₂ to CH₂ can be convincingly understood by examining the composition of the \tilde{a}^1A_1 and \tilde{c}^1A_1 valence complete active space (CAS) wave functions of CH₂:



The “0.98” configuration of the CH₂ \tilde{a}^1A_1 state hinders the approach of N₂ along the C_2 axis, explaining the presence of the energy barrier along this path. On the other hand, the “0.72” configuration of the incoming \tilde{c}^1A_1 state favors bonding along the C_2 axis, rationalizing the actual formation of a C_{2v} geometry stable CH₂N₂ species.⁴ Therefore, even the qualitative description of the diazomethane dissociation would require at least a two-reference wave function, while a single-reference method

would dramatically fail in this respect. For instance, utilization of the “0.98” configuration as a reference wave function should lead to an unbound C_{2v} CH₂N₂ system, while the use of the “0.21” configuration should, if calculable, lead to a barrierless potential energy curve in C_{2v} symmetry. Indeed, Boldyrev et al. mention that no barrier is found at the single reference MP2 level of theory,¹⁵ apparently due to the participation of the “0.72” component in the ground 1A_1 state of CH₂N₂.

In light of the above, we believe that the multireference character of the CH₂N₂ dissociation process along its ground-state PES could provide a good testing ground for the newly developed MRBWCCSD method.

2. Methodology

The MRBWCCSD method is a size-consistent and approximately size-extensive multireference coupled-cluster method based on the use of the Brillouin–Wigner resolvent. In the present problem, the size-nonextensivity error ranges from 20 to 40 μE_h ; thus, the results are practically size-extensive. Details of the MRBWCCSD ansatz have been extensively discussed previously.^{16–20} The Hartree–Fock ground-state electronic configuration of diazomethane in C_s symmetry is

$$\tilde{X}^1A' = (\text{core})^6 (4a')^2 (5a')^2 (6a')^2 (7a')^2 (8a')^2 (1a'')^2 (2a'')^2 (9a')^2 \quad (8)$$

where “core” denotes the total of six 1s-like electrons of C, N and N, respectively. The reference space for the MRBWCCSD calculations contains the (9a') and (10a') orbitals; therefore, it




Figure 1. Potential energy curves of the $\text{CH}_2(\tilde{a}^1A_1)$ approach to $\text{N}_2(X^1\Sigma_g^+)$ for different ϕ angles at CASPT3/cc-pVTZ and MRBWCCSD/cc-pVTZ levels of theory. All energies have been shifted by $+148 E_h$.

is a 2-electron/2-orbital complete active space (CAS) with the resulting CASSCF wave function composed of four singlet




Figure 2. Three-dimensional illustration of the MRBWCCSD potential energy curves of Figure 1.

determinants (three singlet configuration functions). This reference space is adequate to correctly describe the two-configuration character of both \tilde{a} and \tilde{c}^1A_1 states of CH_2 . For purpose of comparison, the PES has also been calculated at the CASPT3 level employing the same CAS reference space, while the MRBWCCSD calculations are performed at the optimized geometries of the aforementioned CASPT3 surface. Note that the present CASPT3 calculation employs a smaller reference space than in ref 4 where a 12-electron/12-orbital reference space was used, the results, however, are practically the same (vide infra). Geometrical and energetical parameters for diazomethane were also calculated at the multireference configuration interaction level (MRCI), as well as the MRCI+Davidson correction (MRCI+Q) and multireference averaged coupled-paired functional (MRACPF) levels of theory. For reasons of uniformity, the cc-pVTZ basis has been employed in all methods.¹⁴ For the MRBWCCSD calculations, we have used a version of the ACES II program extended for MRBWCCSD calculations;^{21,22} the CASPT3, MRCI, MRCI+Q, and MRACPF results were obtained with the MOLPRO 2002 package.²³

3. Results and Discussion

According to the results of Table 1, the first four low-lying states of CH_2 are described satisfactorily in all levels of theory as compared to either experiment or with the very large MRCI/d-aug-cc-pCV6Z calculations of Kalemios et al.²⁴ Note that the “renowned” $\tilde{a}^1A_1 \leftarrow \tilde{X}^3B_1$ energy gap of methylene is better predicted with MRBWCCSD (underestimated by 0.5 kcal/mol) than, for instance, by CASPT3 which, in turn, overestimates it by about 1.5 kcal/mol.

The CASPT3/cc-pVTZ-optimized bond length of N_2 is $r_e = 1.1048 \text{ \AA}$, in very good agreement with the experimental value of 1.0977 \AA .³⁰ The CASPT3 and MRBWCCSD energies at the CASPT3 r_e are -109.36754 and $-109.35775 E_h$, respectively.

Figure 1 depicts different “slices” of the CASPT3 and MRBWCCSD ground-state PES of CH_2N_2 along the $r_{\text{C-N}}$ and ϕ (“attack angle”) coordinates. A three-dimensional representation of the MRBWCCSD curves is illustrated in Figure 2. A detailed explanation of the shape of the surface has already been given,⁴ therefore only the main points will be summarized here. Ultimately, the surface shape is determined by the wave function composition (eq 7) of the \tilde{a}^1A_1 and \tilde{c}^1A_1 states of CH_2 and their relevant energy difference. The largest barrier, but also the

TABLE 2: Geometry and Energetics of CH₂N₂ at Various Multireference Levels of Theory (cc-pVTZ basis)

	CASPT3	CASPT3 ^a	MRCI	MRCI+Q	MRACPF	MRBWCCSD	expt ^b
$E(E_h)$	-148.46979	-148.46935	-148.417234	-148.480411 ^c	-148.482457	-148.47570 ^d	
D_e (kcal/mol) ^e	40.3	38.2	36.6	38.5 ^c	38.6	35.9 ^d	
D_0 (kcal/mol) ^f	23.9	21.6	21.2	23.2 ^c	23.3	21.4 ^d	$\leq 35,^g \leq 44,^h \leq 41.7,^i 25^j$
barrier height (kcal/mol) ^k	34.5	~26	36.2	34.8 ^c	34.4	32.7 ^d	
r_{C-N} (Å)	1.300	1.303	1.295		1.301		1.300
r_{N-N} (Å)	1.129	1.128	1.125		1.139		1.139
r_{C-H} (Å)	1.072	1.071	1.068		1.074		1.077
$\angle HCH$ (deg)	125.4	125.6	125.2		125.1		126.2
ϕ (deg)	0.0	0.0	0.0		0.0		0.0
$\angle NNC$ (deg)	180.0	180.0	180.0		180.0		180.0

^a Reference 4. Basis set: cc-pVTZ, CAS active space: 12 electrons/12 orbitals. ^b Experimental geometrical values from ref 6. ^c Calculated at the MRCI optimized geometry. ^d Calculated at the CASPT3 optimized geometry, see text for details. ^e D_e values with respect to the *adiabatic* CH₂(\tilde{a}^1A_1) + N₂(X¹ Σ_g^+) fragments. ^f D_0 values with respect to the *ground state* CH₂(\tilde{X}^3B_1) + N₂(X¹ Σ_g^+) fragments; see text for details. ^g Pyrolysis, ref 31. ^h Electron impact, ref 32. ⁱ Photodissociation measurements, ref 33. ^j Flash photolysis, ref 34. ^k Barrier height with respect to the CH₂(\tilde{a}^1A_1) + N₂(X¹ Σ_g^+) dissociation limit.

largest depth in the PES is observed at $\phi = 0^\circ$, i.e., along the C₂ axis. The perpendicular attack ($\phi = 90^\circ$) is, as mentioned already, barrierless, but the C–N bond is “weaker” than in any other ϕ orientation. As expected, for intermediate ϕ values (15–75°) the bonding situation becomes a “crossbreed” of the two extreme cases, $\phi = 0^\circ$ and $\phi = 90^\circ$. The participation of the \tilde{c}^1A_1 state of CH₂ becomes evident for $r_{C-N} < 2.3$ Å and is corroborated by the change in wave function composition after the energy maximum, and the sudden opening of the $\angle HCH$ angle from 102 to 126°. Obviously, the involvement of the \tilde{c}^1A_1 state is maximized at $\phi = 0^\circ$.

As shown in Figure 1, the PES morphologies are very similar in both MRBWCCSD and CASPT3 methods. The only qualitative difference between the two methods is that the MRBWCCSD perpendicular ($\phi = 90^\circ$) attack of N₂ to CH₂(\tilde{a}^1A_1) is not predicted to be barrierless, but it exhibits a small barrier of 1.1 kcal/mol. It is possible though that this is an artifact of the calculation since the MRBWCCSD calculations are performed on the CASPT3-optimized geometry.

Relevant numerical data for CH₂N₂ are collected in Table 2. The calculated equilibrium geometry in all levels of theory is in very good agreement with the experimental one. The H₂C–N₂ MRBWCCSD binding energy with respect to the *adiabatic* CH₂(\tilde{a}^1A_1) + N₂(X¹ Σ_g^+) dissociation channel is $D_e = 35.9$ kcal/mol, with the corresponding MRACPF value being 38.6 kcal/mol. Since literature experimental data refer to dissociation to the *ground-state* CH₂(\tilde{X}^3B_1) + N₂(X¹ Σ_g^+) channel, our MRBWCCSD calculated binding energy with respect to the *ground-state* fragments, taking into account the zero-point energy of 5.6 kcal/mol,⁴ is $D_0 = 21.4$ kcal/mol. The corresponding CASPT3, MRCI, MRCI+Q, and MRACPF D_0 values are 21.6, 21.2, 23.2, and 23.3 kcal/mol. There are many different experimental upper bound estimates in the literature regarding the binding energy of the H₂C–N₂ bond.^{31–33} Even though the purpose of this paper is not to extract the most accurate D_e value of CH₂N₂, we note that all our values are in fair agreement with the work of Braun et al., $D_0 = 25$ kcal/mol.³⁴ Finally, all the methods used in the present work predict a barrier to dissociation of 33–35 kcal/mol.

4. Concluding Remarks

The potential energy surface of diazomethane (CH₂N₂) in its ground state along the r_{C-N} and ϕ (“attack” angle of N₂ to CH₂ on the C_s plane) coordinates was studied with the CASPT3 and the MRBWCCSD methods, and energetics were contrasted to the MRCI, MRCI+Q and MRACPF methods. The inherent multireference character of the surface due to the two-

configurational description of the CH₂ \tilde{a}^1A_1 and \tilde{c}^1A_1 states, and the involvement of the latter to the formation of stable diazomethane could adequately be described by the MRBWCCSD method. Guided by the latest successful tests of the method, as well as the very recent inclusion of triple excitations in the MRBWCCSD ansatz,³⁵ we will continue examining its capabilities as a multireference coupled-cluster alternative to conventional approaches for the description of small and medium-sized chemical systems, specifically in cases where single-reference coupled cluster methods are problematic.

Acknowledgment. This work has been supported by the European Community by a Marie Curie Fellowship IHP under Contract No. HPMT-CT-2000-00022. We acknowledge also the support of the COST D23 action (P.C. Grant OC D23.001 from the Ministry of Education, Youth, and Sports of the Czech Republic). The authors also thank Dr. A. Papakondylis for valuable assistance. The computing time provided by the National Center for Scientific Research, DEMOKRITOS, is appreciated.

References and Notes

- Fieser, L. F.; Fieser, M. *Advanced Organic Chemistry*; Reinhold Publishing Corporation: New York, 1961; p 376.
- Bohn, R. B.; Sandford, S. A.; Allamandola, L. J.; Cruikshank, D. P. *Icarus* **1994**, *111*, 151.
- Moore, M. H.; Hudson, R. L. *Icarus* **2003**, *161*, 486 and references therein.
- Papakondylis, A.; Mavridis, A. *J. Phys. Chem. A* **1999**, *103*, 1255 and references therein.
- Kapellos, S. T.; Mavridis, A. *J. Mol. Struct. (THEOCHEM)* **1993**, *279* 151 and references therein.
- Cox, A. P.; Thomas, L. F.; Sheridan, J. *Nature (London)* **1958**, *181*, 1000.
- Pierce, L.; Dobyms, V. *J. Am. Chem. Soc.* **1962**, *84*, 2651.
- See for instance: Streitwieser, A., Jr.; Heathcock, C. H. *Introduction to Organic Chemistry*, 3rd ed.; MacMillan Publishing Company: New York, 1985; p 738.
- For some general thoughts regarding the “resonance” concept, see: Kerkines, I. S. K.; Papakondylis, A.; Mavridis, A. *J. Phys. Chem. A* **2002**, *106*, 4435.
- Walsh, S. P.; Goddard, W. A., III. *J. Am. Chem. Soc.* **1975**, *97*, 5319.
- Gerratt, J.; Cooper, D. L.; Karadakov, P. B.; Raimondi, M. *Chem. Soc. Rev.* **1997**, *87* and references therein.
- Lievín, J.; Verhaegen, G. *Theor. Chim. Acta (Berlin)* **1976**, *42*, 47.
- Lievín, J.; Verhaegen, G. *Theor. Chim. Acta (Berlin)* **1977**, *45*, 269.
- Dunning, T. H., Jr. *J. Chem. Phys.* **1989**, *90*, 1007.
- Boldyrev, A. I.; Schleyer, P. v. R.; Higgins, D.; Thomson, C.; Kramarenko, S. S. *J. Comput. Chem.* **1992**, *13*, 1066.
- Čársky, P.; Hrouda, V.; Sychrovský, V.; Hubač, I.; Babinec, P.; Mach, P.; Urban, J.; Mášik, J. *Collect. Czech. Chem. Commun.* **1995**, *60*, 1419.
- Mášik, J.; Hubač, I. In *Quantum Systems in Chemistry and Physics: Trends in Methods and Applications*; McWeeny, R., Maruani, J.,

Smeyers, Y. G.; Wilson, S., Eds.; Kluwer Academic: Dordrecht, The Netherlands, 1997; pp 283–308 and references therein.

(18) Kerkines, I. S. K.; Pittner, J.; Čársky, P.; Mavridis, A.; Hubač, I. *J. Chem. Phys.* **2002**, *117*, 9733.

(19) Teberekidis, V. I.; Kerkines, I. S. K.; Tsipis, C. A.; Čársky, P.; Mavridis, A.; *Int. J. Quantum Chem.* **2005**, *102*, 762.

(20) Kardahakis, S.; Pittner, J.; Čársky, P.; Mavridis, A. *Int. J. Quantum Chem.* **2005**, *104*, 458.

(21) ACES II is a program product of Quantum Theory Project, University of Florida: Stanton, J. F.; Gauss, J.; Watts, J. D.; Nooijen, M.; Oliphant, N.; Perera, S. A.; Szalay, P. G.; Lauderdale, W. J.; Gwaltney, S. R.; Beck, S.; Balková, A.; Bernholdt, D. E.; Baeck, K.-K.; Rozyczko, P.; Sekino, H.; Huber, C.; Bartlett, R. J. Integral packages included are vmol (Almlöf, J.; Taylor, P. R.); vprops (Taylor, P. R.); and abacus (Helgaker, T.; Jensen, H. J. Aa.; Jørgensen, P.; Olsen, J.; Taylor, P. R.).

(22) Pittner, J.; Nachtigall, P.; Čársky, P.; Mášik, J.; Hubač, I. *J. Chem. Phys.* **1999**, *110*, 10275.

(23) MOLPRO a package of ab initio programs designed by: Werner, H.-J.; Knowles, P. J. Version 2002.6: Amos, R. D.; Bernhardsson, A.; Berning, A.; Celani, P.; Cooper, D. L.; Deegan, M. J. O.; Dobbyn, A. J.; Eckert, F.; Hampel, C.; Hetzer, G.; Knowles, P. J.; Korona, T.; Lindh, R.;

Lloyd, A. W.; McNicholas, S. J.; Manby, F. R.; Meyer, W.; Mura, M. E.; Nicklass, A.; Palmieri, P.; Pitzer, R.; Rauhut, G.; Schütz, M.; Schumann, U.; Stoll, H.; Stone, A. J.; Tarroni, R.; Thorsteinsson, T.; Werner, H.-J.

(24) Kalemos, A.; Dunning, T. H., Jr.; Mavridis, A.; Harrison, J. F. *Can. J. Chem.* **2004**, *82*, 684.

(25) Jensen, P.; Bunker, P. R. *J. Chem. Phys.* **1988**, *89*, 1327.

(26) Czászár, A. G.; Leininger, M. L.; Szalay, V. *J. Chem. Phys.* **2003**, *118*, 10631.

(27) Petek, H.; Nesbitt, D. J.; Darwin, D. C.; Ogilby, P. R.; Moore, C. B.; Ramsay, D. A. *J. Chem. Phys.* **1989**, *91*, 6566.

(28) Duxbury, G.; Jungen, Ch. *Mol. Phys.* **1988**, *63*, 981.

(29) Herzberg, G.; Johns, J. W. C. *Proc. R. Soc. London A* **1966**, *295*, 107.

(30) Huber, K. P.; Herzberg, G. H. *Molecular Spectra and Molecular Structure, Vol. IV: Constants of Diatomic Molecules*; Van Nostrand Reinhold: New York, 1979.

(31) Setzer, D. W.; Rabinovich, B. S. *Can. J. Chem.* **1962**, *40*, 1425.

(32) Paulett, G. S.; Ettinger, P. E. *J. Chem. Phys.* **1963**, *39*, 3534.

(33) Laufer, A. H.; Okabe, H. *J. Am. Chem. Soc.* **1971**, *93*, 4137.

(34) Braun, W.; Bass, A. M.; Pilling, M. *J. Chem. Phys.* **1970**, *52*, 5131.

(35) Pittner, J.; Demel, O. *J. Chem. Phys.* **2005**, *122*, 181101.