

## Exact solution of the Schrodinger equation for a particle in a tetrahedral box

This content has been downloaded from IOPscience. Please scroll down to see the full text.

1982 J. Phys. A: Math. Gen. 15 2131

(<http://iopscience.iop.org/0305-4470/15/7/024>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

### Download details:

IP Address: 195.134.76.74

This content was downloaded on 01/09/2016 at 10:35

Please note that [terms and conditions apply](#).

You may also be interested in:

[On the quantum particle in a polyhedral box](#)

J W Turner

[A Lie group approach to the Schrodinger equation for a particle in an equilateral triangular infinite well](#)

Wajdi Gaddah

[Exact solution of the particle-rigid-rotor scattering problem](#)

V N Ostrovsky and V I Ustimov

[Normal electric modes at small polyhedral particles](#)

D Langbein

[Electromagnetic selection rules in the triangular -cluster model of  \$^{12}\text{C}\$](#)

G Stellin, L Fortunato and A Vitturi

[Group theoretical treatments of strongly correlated atomic dynamics](#)

A R P Rau

[A lower bound for ground-state energy by Steiner symmetrisation of the potential](#)

R Subramanian and K V Bhagwat

## Exact solution of the Schrödinger equation for a particle in a tetrahedral box

H R Krishnamurthy†, H S Mani and H C Verma‡

Indian Institute of Technology, Kanpur-208016, India

Received 8 December 1981, in final form 25 February 1982

**Abstract.** We obtain the exact solution of the Schrödinger equation for a particle confined to (i) an equilateral triangle, (ii) a tetrahedral box with corners  $(-\pi/\sqrt{2}, -\pi/\sqrt{2}, -\pi/2)$ ,  $(\pi/\sqrt{2}, -\pi/\sqrt{2}, \pi/2)$ ;  $(-\pi/\sqrt{2}, \pi/\sqrt{2}, \pi/2)$  and  $(\pi/\sqrt{2}, \pi/\sqrt{2}, -\pi/2)$ . The energies are: for (i),  $E_{nl} = (8\hbar^2\pi^2/9mL^2)(n^2 + l^2 - nl)$  where  $L$  is the side of the triangle and  $l, n$  are distinct non-zero integers and for (ii),  $E_{lmn} = (\hbar^2/8m) \times [3(l^2 + m^2 + n^2) - 2lm - 2mn - 2nl]$  where  $l, m$  and  $n$  are distinct non-zero integers. The wavefunctions have been classified according to the irreducible representation of the corresponding symmetry groups.

### 1. Introduction

Exact solutions of the Schrödinger equation for a particle confined to certain regions of space (either two-dimensional or three-dimensional) have been found only for a few cases. We have found an amusing way to obtain the solution of the Schrödinger equation for: (i) a particle confined to an equilateral triangle in two dimensions; (ii) a particle confined in a tetrahedral box whose corners are  $(-\pi/\sqrt{2}, -\pi/\sqrt{2}, -\pi/2)$ ,  $(\pi/\sqrt{2}, -\pi/\sqrt{2}, \pi/2)$ ;  $(-\pi/\sqrt{2}, \pi/\sqrt{2}, \pi/2)$  and  $(\pi/\sqrt{2}, \pi/\sqrt{2}, -\pi/2)$ . Though (i) has been solved before using other methods, we believe that the results for the tetrahedral box are new. The solution to the problem is based on the Schrödinger equation of  $N$  hard cores confined to a length  $L$  in one dimension, the exact solution of which is known (Lieb and Mattis 1966). It is possible to transform the  $N = 3$  and 4 cases of the hard-core problem into the problem of a single particle confined to a region in two and three dimensions respectively. It turns out that the region of confinement is an equilateral triangle for the two-dimensional case and a tetrahedron with corners specified as above for the case of three dimensions. To make the article self-contained, we reproduce the essential results of the  $N$ -hard-core problem in § 2. In § 3 we illustrate our method of transforming the equation of  $N$  hard cores to the Schrödinger equation of a single particle for the  $N = 3$  case in detail. We then quote in § 4 the results for the  $N = 4$  case. The wavefunctions have been classified according to the irreducible representations of the respective symmetry groups.

† Department of Physics, Indian Institute of Science, Bangalore 560012, India.

‡ Department of Physics, Patna Science College, Patna 800005, India.

## 2. Problem of $N$ hard cores in one dimension

Consider  $N$  hard cores (molecules of vanishingly small length and mass  $m$ ) whose coordinates  $x_i$  are confined to lie between 0 and  $L$ ,

$$0 \leq x_i \leq L \quad i = 1, 2, \dots, n. \quad (2.1)$$

The cores are free to move in the allowed region and are to be treated as identical fermions. In particular, the wavefunction

$$\varphi(x_1 \dots x_N) = 0 \quad (2.2)$$

whenever

$$x_i = x_j, \quad i, j = 1, \dots, N.$$

The Schrödinger equation

$$-\frac{\hbar^2}{2m} \sum_i \frac{\partial^2 \varphi}{\partial x_i^2} = E\varphi \quad (2.3)$$

with the above conditions (2.1) and (2.2), plus rigid boundary conditions has the following energy eigenvalues (Lieb and Mattis 1966):

$$En_1 \dots n_N = (\hbar^2 \pi^2 / 2mL^2)(n_1^2 + n_2^2 + \dots + n_N^2) \quad (2.4)$$

where  $n_i$  are distinct positive integers. The corresponding wavefunctions are the determinants

$$\varphi_{n_1 \dots n_N}(x_1 \dots x_N) = \begin{vmatrix} \sin k_1 x_1 & \sin k_1 x_2 & \dots & \sin k_1 x_N \\ \vdots & \vdots & & \vdots \\ \sin k_N x_1 & \sin k_N x_2 & \dots & \sin k_N x_N \end{vmatrix} \quad (2.5)$$

where  $k_i = \pi n_i / L$ .

We can state the same problem in terms of periodic boundary conditions, which is the form used in the present investigation. In this case the wavefunction obeys the boundary conditions (using coordinates  $\theta_i$  in place of  $x_i$ ):

$$\varphi(\theta_1, \dots, \theta_j + 2\pi, \dots, \theta_N) = \varphi(\theta_1, \dots, \theta_j, \dots, \theta_N) \quad j = 1, \dots, N \quad (2.6)$$

For these boundary conditions, the energy levels are

$$En_1 \dots n_N = \frac{\hbar^2}{2m}(n_1^2 + n_2^2 + \dots + n_N^2) \quad (2.7)$$

and the wavefunction is the determinant

$$\varphi_{n_1 \dots n_N}(\theta_1 \dots \theta_N) = \begin{vmatrix} \exp(in_1 \theta_1) & \dots & \exp(in_1 \theta_N) \\ \exp(in_2 \theta_1) & & \exp(in_2 \theta_N) \\ \vdots & & \vdots \\ \exp(in_N \theta_1) & & \exp(in_N \theta_N) \end{vmatrix}. \quad (2.8)$$

### 3. Particle in an equilateral triangle

A useful transformation of the variables  $\theta_1 \dots \theta_N$  which allows us to separate the centre of mass of the whole system is

$$\begin{aligned}
 z_1 &= \theta_1 - \theta_2 \\
 z_2 &= \frac{1}{2}(\theta_1 + \theta_2) - \theta_3 \\
 &\vdots \\
 z_i &= i^{-1}(\theta_1 + \dots + \theta_i) - \theta_{i+1}, \quad i \leq N-1 \\
 &\vdots \\
 z_N &= N^{-1}(\theta_1 + \dots + \theta_N).
 \end{aligned}
 \tag{3.1}$$

The Schrödinger equation in terms of the new variables is

$$-\frac{\hbar^2}{2m} \left[ 2 \frac{\partial^2}{\partial z_1^2} + \dots + \frac{i+1}{i} \frac{\partial^2}{\partial z_i^2} + \dots + \frac{N}{(N-1)} \frac{\partial^2}{\partial z_{N-1}^2} + \frac{1}{N} \frac{\partial^2}{\partial z_N^2} \right] \varphi = E\varphi.
 \tag{3.2}$$

For  $N = 3$  these become

$$z_1 = \theta_1 - \theta_2 \quad z_2 = \frac{1}{2}(\theta_1 + \theta_2) - \theta_3 \quad z_3 = \frac{1}{3}(\theta_1 + \theta_2 + \theta_3)
 \tag{3.3}$$

and

$$-\frac{\hbar^2}{2m} \left[ 2 \frac{\partial^2}{\partial z_1^2} + \frac{3}{2} \frac{\partial^2}{\partial z_2^2} + \frac{1}{3} \frac{\partial^2}{\partial z_3^2} \right] \varphi = E\varphi.
 \tag{3.4}$$

The  $z_3$  coordinate describes motion associated with the centre of mass and the dependence of  $\varphi$  on it is separable. If we think of  $(z_1/\sqrt{2})$  and  $z_2\sqrt{2}/\sqrt{3}$  as the  $x$  and the  $y$  coordinates of a single particle, then we reproduce the Schrödinger equation for a single particle. However, the boundary conditions which were simple in terms of  $\theta$ 's have to be transformed in terms of new variables. Using

$$\begin{aligned}
 Y_1 &= z_1/\sqrt{2} = \frac{1}{2}(\theta_1 - \theta_2) \\
 Y_2 &= z_2\sqrt{2}/\sqrt{3} = \sqrt{\frac{2}{3}}[\frac{1}{2}(\theta_1 + \theta_2) - \theta_3] \\
 Y_3 &= \sqrt{3}z_3 = (\theta_1 + \theta_2 + \theta_3)/\sqrt{3}.
 \end{aligned}
 \tag{3.5}$$

We can find the restriction on the domain of the  $Y_1, Y_2$  plane. To achieve this we invert equation (3.5) and get

$$\begin{aligned}
 \theta_1 &= Y_1/\sqrt{2} + Y_2/\sqrt{6} + Y_3/\sqrt{3} \\
 \theta_2 &= -Y_1/\sqrt{2} + Y_2/\sqrt{6} + Y_3/\sqrt{3} \\
 \theta_3 &= -2Y_2/\sqrt{6} + Y_3/\sqrt{3}.
 \end{aligned}
 \tag{3.6}$$

Without loss of generality, we can restrict the  $\theta$ -space to the region  $\theta_1 \geq \theta_2, \theta_2 \geq \theta_3$  and  $\theta_3 + 2\pi \geq \theta_1$ . In terms of the  $Y$ 's these translate into the restrictions

$$Y_1 \geq 0, \quad Y_1 \leq \sqrt{3} Y_2, \quad Y_1/\sqrt{2} \leq 2\pi - Y_2\sqrt{\frac{3}{2}}
 \tag{3.7}$$

respectively. This corresponds to the particle being confined in an equilateral triangle with corners  $(0, 0), (\sqrt{2}\pi, \sqrt{\frac{2}{3}}\pi)$  and  $(0, 2\sqrt{\frac{2}{3}}\pi)$ . The wavefunction for the single

particle can be obtained from equation (2.8) using the transformations (3.6).

$$\begin{aligned}
 \varphi(Y_1, Y_2, Y_3) &= \\
 & \left. \begin{matrix} n_1, n_2, n_3 \\ \exp\left[in_1\left(\frac{Y_1}{\sqrt{2}} + \frac{Y_2}{\sqrt{6}} + \frac{Y_3}{\sqrt{3}}\right)\right] & \exp\left[in_1\left(-\frac{Y_1}{\sqrt{2}} + \frac{Y_2}{\sqrt{6}} + \frac{Y_3}{\sqrt{3}}\right)\right] & \exp\left[+in_1\left(-\frac{2Y_2}{\sqrt{6}} + \frac{Y_3}{\sqrt{3}}\right)\right] \\ \exp\left[in_2\left(\frac{Y_1}{\sqrt{2}} + \frac{Y_2}{\sqrt{6}} + \frac{Y_3}{\sqrt{3}}\right)\right] & \exp\left[in_2\left(-\frac{Y_1}{\sqrt{2}} + \frac{Y_2}{\sqrt{6}} + \frac{Y_3}{\sqrt{3}}\right)\right] & \exp\left[in_2\left(-\frac{2Y_2}{\sqrt{6}} + \frac{Y_3}{\sqrt{3}}\right)\right] \\ \exp\left[in_3\left(\frac{Y_1}{\sqrt{2}} + \frac{Y_2}{\sqrt{6}} + \frac{Y_3}{\sqrt{3}}\right)\right] & \exp\left[in_3\left(-\frac{Y_1}{\sqrt{2}} + \frac{Y_2}{\sqrt{6}} + \frac{Y_3}{\sqrt{3}}\right)\right] & \exp\left[in_3\left(-\frac{2Y_2}{\sqrt{6}} + \frac{Y_3}{\sqrt{3}}\right)\right] \end{matrix} \right| \\
 & = \exp[i(n_1 + n_2 + n_3)Y_3/\sqrt{3}] \\
 & \times \left. \begin{matrix} 1 & 1 & 1 \\ \exp\left[i(n_2 - n_1)\left(\frac{Y_1}{\sqrt{2}} + \frac{Y_2}{\sqrt{6}}\right)\right] & \exp\left[i(n_2 - n_1)\left(-\frac{Y_1}{\sqrt{2}} + \frac{Y_2}{\sqrt{6}}\right)\right] & \exp\left[i(n_2 - n_1)\left(-\frac{2Y_2}{\sqrt{6}}\right)\right] \\ \exp\left[i(n_3 - n_1)\left(\frac{Y_1}{\sqrt{2}} + \frac{Y_2}{\sqrt{6}}\right)\right] & \exp\left[i(n_3 - n_1)\left(-\frac{Y_1}{\sqrt{2}} + \frac{Y_2}{\sqrt{6}}\right)\right] & \exp\left[i(n_3 - n_1)\left(-\frac{2Y_2}{\sqrt{6}}\right)\right] \end{matrix} \right| \tag{3.8}
 \end{aligned}$$

The factor  $\exp[i(n_1 + n_2 + n_3)Y_3/\sqrt{3}]$  will be suppressed as it is separable and refers to the centre of mass motion of the three hard cores. We are interested only in the  $Y_1, Y_2$  part. Writing

$$\phi_{l,m}(Y_1, Y_2) = \left. \begin{matrix} 1 & 1 & 1 \\ \exp\left[il\left(\frac{Y_1}{\sqrt{2}} + \frac{Y_2}{\sqrt{6}}\right)\right] & \exp\left[il\left(-\frac{Y_1}{\sqrt{2}} + \frac{Y_2}{\sqrt{6}}\right)\right] & \exp\left[il\left(-\frac{2Y_2}{\sqrt{6}}\right)\right] \\ \exp\left[im\left(\frac{Y_1}{\sqrt{2}} + \frac{Y_2}{\sqrt{6}}\right)\right] & \exp\left[im\left(-\frac{Y_1}{\sqrt{2}} + \frac{Y_2}{\sqrt{6}}\right)\right] & \exp\left[im\left(-\frac{2Y_2}{\sqrt{6}}\right)\right] \end{matrix} \right| \tag{3.9}$$

we immediately see that it satisfies the Schrödinger equation

$$-(\hbar^2/2m)[\partial^2/\partial Y_1^2 + \partial^2/\partial Y_2^2]\phi_{l,m}(Y_1, Y_2) = E_{l,m}\phi_{l,m}(Y_1, Y_2) \tag{3.10}$$

with

$$E_{l,m} = (\hbar^2/3m)(l^2 + m^2 - lm). \tag{3.11}$$

This can either be obtained from

$$E_{n_1, n_2, n_3} = (\hbar^2/2m)(n_1^2 + n_2^2 + n_3^2) = (\hbar^2/2m)[\frac{1}{3}(n_1 + n_2 + n_3)^2 + \frac{2}{3}(l^2 + m^2 - lm)]$$

and removing the centre of mass energy or by direct substitution of  $\phi_{l,m}$  from equation (3.9) in equation (3.10).

For an equilateral triangle of side  $L$ , we will have the energy as

$$E_{l,m} = (8\hbar^2 \pi^2/9mL^2)(l^2 + m^2 - lm) \tag{3.12}$$

and  $Y_i$  ( $i = 1, 2$ ) are to be interpreted as  $(2\pi\sqrt{\frac{2}{3}}Y_i/L)$  in equation (3.9). This has been derived using other methods (Lame 1852, Mathews and Walker 1970).

We now discuss the symmetry of the solutions. It is clear that we have six symmetry operations under which the Schrödinger equation with the boundary conditions remains invariant, which form the group  $\sigma_{3v}$ .

We classify our solutions in terms of the irreducible representations  $A_1$ ,  $A_2$  and  $E$  of  $\sigma_{3v}$  (Landau and Lifshitz 1958). In order to do this it is more convenient to transform the  $Y_1, Y_2$  coordinates so that the origin coincides with the centre of the triangle. We use

$$Y_1 = y_1 + \sqrt{2}\pi/3, \quad Y_2 = y_2 + \sqrt{\frac{2}{3}}\pi \tag{3.13}$$

and write

$$\phi_{lm}(y_1, y_2) = \begin{vmatrix} 1 & 1 & 1 \\ \exp\left[il\left(\frac{y_1}{\sqrt{2}} + \frac{y_2}{\sqrt{6}} + \frac{2\pi}{3}\right)\right] & \exp\left[il\left(-\frac{y_1}{\sqrt{2}} + \frac{y_2}{\sqrt{6}}\right)\right] & \exp\left[-il\left(\frac{2y_2}{\sqrt{6}} + \frac{2\pi}{3}\right)\right] \\ \exp\left[im\left(\frac{y_1}{\sqrt{2}} + \frac{y_2}{\sqrt{6}} + \frac{2\pi}{3}\right)\right] & \exp\left[im\left(-\frac{y_1}{\sqrt{2}} + \frac{y_2}{\sqrt{6}}\right)\right] & \exp\left[-im\left(\frac{2y_2}{\sqrt{6}} + \frac{2\pi}{3}\right)\right] \end{vmatrix} \tag{3.14}$$

The operation of  $C_3$  and  $\sigma_v$  in terms of  $y_1$  and  $y_2$  are

$$C_3: \quad y_1 \rightarrow -\frac{1}{2}(y_1 - \sqrt{3}y_2) \quad y_2 \rightarrow -\frac{1}{2}(\sqrt{3}y_1 + y_2) \tag{3.15}$$

and

$$\sigma_v: \quad y_1 \rightarrow y_1 \quad y_2 \rightarrow -y_2.$$

It is easy to verify that

$$C_3\phi_{lm} = \exp[i\frac{2}{3}\pi(l+m)]\phi_{lm}$$

and

$$\sigma_v\phi_{lm} = -\exp[i\frac{2}{3}\pi(l+m)]\phi_{lm}^*. \tag{3.16}$$

It immediately follows that, if  $l+m = 3p$  where  $p$  is an integer

$$\frac{1}{2}(\phi_{lm} - \phi_{lm}^*) \text{ transforms as } A_1 \text{ and}$$

$$\frac{1}{2}(\phi_{lm} + \phi_{lm}^*) \text{ transforms as } A_2.$$

On the other hand, if  $l+m = 3p+1$  or  $3p+2$ ,  $\phi_{lm}$  and  $\phi_{l,m}^*$  form a basis for the irreducible representation  $E$ .

#### 4. Particle in a tetrahedron

The procedure which leads to the solution of a particle in a tetrahedron with corners A  $(-\pi/\sqrt{2}, -\pi/\sqrt{2}, -\pi/2)$ ; B  $(\pi/\sqrt{2}, -\pi/\sqrt{2}, \pi/2)$ ; C  $(-\pi/\sqrt{2}, \pi/\sqrt{2}, \pi/2)$  and D  $(\pi/\sqrt{2}, \pi/\sqrt{2}, -\pi/2)$  is similar to the one described in the previous section. Here we work with the  $N = 4$  solution of equation (2.8). We will not repeat all the details, but only write down some of the relevant steps and the final results. The transformations are

$$\begin{aligned} \theta_1 &= \frac{1}{2}y_4 + \frac{1}{2}y_3 + \frac{1}{2}y_1\sqrt{2} + \frac{3}{4}\pi & \theta_2 &= \frac{1}{2}y_4 - \frac{1}{2}y_3 + \frac{1}{2}y_2\sqrt{2} + \frac{1}{4}\pi \\ \theta_3 &= \frac{1}{2}y_4 + \frac{1}{2}y_3 - \frac{1}{2}y_1\sqrt{2} - \frac{1}{4}\pi & \theta_4 &= \frac{1}{2}y_4 - \frac{1}{2}y_3 - \frac{1}{2}y_2\sqrt{2} - \frac{3}{4}\pi. \end{aligned} \tag{4.1}$$

The conditions  $\theta_1 \geq \theta_2$ ,  $\theta_2 \geq \theta_3$ ,  $\theta_3 \geq \theta_4$  and  $\theta_4 + 2\pi \geq \theta_1$  lead to the particle being confined in the tetrahedron ABCD in the  $y_1, y_2, y_3$  space. The wavefunction after removing the centre of mass coordinate is the  $4 \times 4$  determinant


$$\phi_{lmn}(y_1, y_2, y_3) = \begin{vmatrix} 1 & 1 & 1 & 1 \\ \exp\left[il\left(\frac{y_1}{\sqrt{2}} + \frac{y_3}{2} + \frac{3\pi}{4}\right)\right] & \exp\left[il\left(\frac{y_2}{\sqrt{2}} - \frac{y_3}{2} + \frac{\pi}{4}\right)\right] & \exp\left[il\left(-\frac{y_1}{\sqrt{2}} + \frac{y_3}{2} - \frac{\pi}{4}\right)\right] & \exp\left[il\left(-\frac{y_2}{\sqrt{2}} - \frac{y_3}{2} - \frac{3\pi}{4}\right)\right] \\ \exp\left[im\left(\frac{y_1}{\sqrt{2}} + \frac{y_3}{2} + \frac{3\pi}{4}\right)\right] & \exp\left[im\left(\frac{y_2}{\sqrt{2}} - \frac{y_3}{2} + \frac{\pi}{4}\right)\right] & \exp\left[im\left(-\frac{y_1}{\sqrt{2}} + \frac{y_3}{2} - \frac{\pi}{4}\right)\right] & \exp\left[im\left(-\frac{y_2}{\sqrt{2}} - \frac{y_3}{2} - \frac{3\pi}{4}\right)\right] \\ \exp\left[in\left(\frac{y_1}{\sqrt{2}} + \frac{y_3}{2} + \frac{3\pi}{4}\right)\right] & \exp\left[in\left(\frac{y_2}{\sqrt{2}} - \frac{y_3}{2} + \frac{\pi}{4}\right)\right] & \exp\left[in\left(-\frac{y_1}{\sqrt{2}} + \frac{y_3}{2} - \frac{\pi}{4}\right)\right] & \exp\left[in\left(-\frac{y_2}{\sqrt{2}} - \frac{y_3}{2} - \frac{3\pi}{4}\right)\right] \end{vmatrix} \quad (4.2)$$

the energy is

$$E_{lmn} = (\hbar^2/8m)[3(l^2 + m^2 + n^2) - 2lm - 2ln - 2mn]. \quad (4.3)$$

The corners of the tetrahedron have been chosen so that the line joining the mid points of AC and BD is the  $y_1$  axis, that joining the mid points of AB and CD is the  $y_2$  axis, and that joining the mid points of AD and BC is the  $y_3$  axis. The tetrahedron goes into itself (see figure 1) under:

- (i) Identity ( $O_1$ ),
- (ii) rotation of  $\pi$  about  $y_1$  axis ( $O_2$ ),
- (iii) rotation of  $\pi$  about  $y_2$  axis ( $O_3$ ),
- (iv) rotation of  $\pi$  about  $y_3$  axis ( $O_4$ ),
- (v) reflection in the plane  $y_1 = y_2$  ( $O_5$ ),
- (vi) reflection in the plane  $y_1 + y_2 = 0$  ( $O_6$ ),
- (vii) rotation of  $\frac{1}{2}\pi$  about the  $y_3$  axis followed by reflection in the plane  $y_3 = 0$  ( $O_7$ ),
- (viii) rotation of  $-\frac{1}{2}\pi$  about the  $y_3$  axis followed by reflection in the plane  $y_3 = 0$  ( $O_8$ ).



**Figure 1.** Geometry of the tetrahedral box for which (4.2) gives the eigenfunctions of the Schrödinger equation.

The operations form an eight-element group  $D_{2d}$ . The irreducible representations are classified as  $A_1$ ,  $A_2$ ,  $B_1$ ,  $B_2$  and  $E$  (Landau and Lifshitz 1958). One also has

$$\begin{aligned} O_2\phi_{lmn} &= -\exp[i(l+m+n)\frac{1}{2}\pi]\phi_{lmn}^* \\ O_3\phi_{lmn} &= -\exp[-i(l+m+n)\frac{1}{2}\pi]\phi_{lmn}^* \\ O_4\phi_{lmn} &= \exp[i(l+m+n)\pi]\phi_{lmn} \\ O_5\phi_{lmn} &= \phi_{lmn}^* \\ O_6\phi_{lmn} &= \exp[i(l+m+n)\pi]\phi_{lmn}^* \\ O_7\phi_{lmn} &= -\exp[i(l+m+n)\frac{1}{2}\pi]\phi_{lmn} \\ O_8\phi_{lmn} &= -\exp[-i(l+m+n)\frac{1}{2}\pi]\phi_{lmn}. \end{aligned}$$

These can be used to determine the symmetry properties of the wavefunctions. In particular,

- (i) for  $l+m+n=4p$ , where  $p$  is an integer, the wavefunction  $(\phi_{l,m,n} - \phi_{l,m,n}^*)$  belongs to the representation  $B_1$  and  $(\phi_{l,m,n} + \phi_{l,m,n}^*)$  belongs to  $B_2$ ;
- (ii) for  $l+m+n=4p+1$ ,  $4p+3$ ;  $\phi_{l,m,n}$  and  $\phi_{l,m,n}^*$  form the basis for the two-dimensional representation  $E$ ; and finally
- (iii) for  $l+m+n=4p+2$ ;  $(\phi_{lmn} + \phi_{lmn}^*)$  belongs to  $A_1$  and  $(\phi_{l,m,n} - \phi_{l,m,n}^*)$  belongs to the representation  $A_2$ .

### Acknowledgment

We would like to thank Dr K Banerjee for reading the manuscript.

### References

- Lame M G 1852 *Leçons sur le Théorie Mathématique d'Elasticité des Cores Solides* (Paris: Bachelier)  
 Landau L D and Lifshitz E M 1958 *Quantum Mechanics* (Oxford: Pergamon) ch XII  
 Lieb E H and Mattis D C 1966 *Mathematical Physics in One Dimension* (New York: Academic) p 396  
 Mathews J and Walker R L 1970 *Mathematical Methods for Physicists* (New York: Benjamin) p 237-9